

Forbidden triples for hamiltonicity

Jan Brousek¹

*Department of Mathematics, University of West Bohemia, P.O. Box 314, 306 14 Plzeň ,
Czech Republic*

Received 9 September 1999; revised 13 June 2000; accepted 3 June 2001

Abstract

In this paper we characterize all triples of connected graphs C, X, Y (where C is a claw) and such that every 2-connected CXY -free graph G is hamiltonian. This result together with a previous result by Faudree, Gould, Jacobson, and Lesniak give a full characterization of triples of forbidden subgraphs implying hamiltonicity of 2-connected graphs. © 2002 Elsevier Science B.V. All rights reserved.

Keywords: Hamiltonian graph; Forbidden subgraph; Claw-free graph

1. Introduction

Throughout the paper, by a *graph* we always mean a finite undirected graph $G = (V(G), E(G))$ without loops and multiple edges. We follow the most common graph-theoretical terminology and notation and for concepts not defined here we refer to [2].

If $M \subset V(G)$ then $\langle M \rangle$ denotes the induced subgraph on M . If $\emptyset \subsetneq M \subsetneq V(G)$ we say that $\langle M \rangle$ is a *proper induced subgraph* of G . If H_1, \dots, H_k are graphs, then we say that a graph G is $H_1 \dots H_k$ -free if G does not contain a copy of any of the graphs H_1, \dots, H_k as an induced subgraph. The graphs H_1, \dots, H_k will be referred to in this context as *forbidden subgraphs*.

For some graphs that will often occur as forbidden subgraphs we will use the fixed terminology and notation, see Fig. 1.

We define \mathcal{P} to be the class of graphs obtained by taking two vertex-disjoint triangles $\langle \{v_1, v_2, v_3\} \rangle, \langle \{w_1, w_2, w_3\} \rangle$ and by joining every pair of vertices $\{v_i, w_i\}$ by a path P_n for $n \geq 3$ or by a triangle. We denote graphs from the class \mathcal{P} by P_{x_1, x_2, x_3} , where $x_i = n$ if v_i, w_i are joined by P_n , and $x_i = T$ if v_i, w_i are joined by a triangle. Graphs from \mathcal{P}

¹ Research supported by grant GA ČR No. 201/97/0407.

E-mail address: brousek@kma.zcu.cz (J. Brousek).




Fig. 1.




Fig. 2.

play an important role throughout the paper. The *racket* *R* is the graph obtained by removing one vertex of degree four from a copy of $P_{T,T,T}$ (see Fig. 2).

The following result was proved in [3].

Theorem A (Brousek [3]). *Every 2-connected non-hamiltonian claw-free graph contains a graph from the class \mathcal{P} as an induced subgraph.*

In [3] are also given as a corollary of Theorem A the following examples of graphs *X*, *Y* such that every 2-connected *CXY*-free graph is hamiltonian.

Theorem B (Brousek [3]). *Let *G* be a 2-connected graph satisfying one of the following conditions.*

- (i) G is $CDP_{3,3,3}$ -free,
- (ii) G is $CEP_{T,T,T}$ -free,
- (iii) G is $CP_7P_{T,T,T}$ -free.

Then G is hamiltonian.

Bedrossian [1] characterized all pairs of forbidden graphs which imply hamiltonicity.

Theorem C (Bedrossian [1]). *Let X and Y be connected graphs, both different from P_3 , and let G be a 2-connected graph that is not a cycle. Then G being XY -free implies G is hamiltonian if and only if (up to a symmetry) X is a claw and $Y \in \{P_4, P_5, P_6, C_3, Z_1, Z_2, B, N, W\}$.*

Faudree et al. in [4] characterized all triples of graphs X, Y, Z such that none of X, Y, Z is isomorphic to the claw and G being 2-connected and XYZ -free implies hamiltonicity.

Theorem D (Faudree et al. [4]). *Let G be a 2-connected graph of sufficient large order and let X, Y, Z be connected graphs different from P_3 and the claw. Then G being XYZ -free implies G is hamiltonian if and only if X, Y, Z are induced subgraphs of one of the following triples:*

- (i) $P_4, K_4 \setminus e, K_{\lfloor n/2 \rfloor + 1}$,
- (ii) $P_5, K_4 \setminus e, K_{2,3}$,
- (iii) $P_6, K_3, K_{2,2}$,
- (iv) $C_{1,2,2}, K_3, K_{2,2}$,
- (v) $P_4, K_{1,1,3}, K_{2,3}$,
- (vi) $P_5, K_3, K_{2, \lfloor n/3 \rfloor}$,
- (vii) $C_{1,1,3}, K_3, K_{2,2}$,
- (viii) $C_{1,1,2}, K_3, K_{2, \lfloor (n-1)/2 \rfloor - 2}$,

where $C_{i,j,k}$ denotes a graph consist of three paths of length $i+1, j+1, k+1$ with one common endvertex.

In the main result of this paper, Theorem 1, we characterize all triples of forbidden subgraphs X, Y, Z which imply hamiltonicity and one of them is the claw. This result together with Theorem D gives a full characterization of forbidden triples of subgraphs for hamiltonicity.

2. Main result

For convenience we will say that a triple of graphs X, Y, Z is a *good triple* if every 2-connected XYZ -free graph is necessarily hamiltonian. We will say a good triple




Fig. 3.

X_1, X_2, X_3 is *maximal* if there is no good triple Y_1, Y_2, Y_3 such that $X_i \subset Y_i$ for $i=1, 2, 3$ (in the sense of induced subgraphs) and $X_i \neq Y_i$ for at least one $i \in \{1, 2, 3\}$ (or, equivalently, if there is no good triple Y_1, Y_2, Y_3 such that the class of $X_1X_2X_3$ -free graphs is a proper subclass of the class of $Y_1Y_2Y_3$ -free graphs). Remark, that with respect to Theorem C we will consider that no forbidden subgraph is an induced subgraph of P_6 , N or W .

Theorem 1. Let X, Y be connected graphs such that neither X nor Y is an induced subgraph of any of the graphs P_6 , N or W . Then

- (i) There exist 62 pairs X, Y of graphs such that the property “ G is 2-connected and CXY -free” implies G is hamiltonian.
- (ii) C, X, Y is a maximal triple of forbidden subgraphs such that G being CXY -free implies G is hamiltonian if and only if $CXY \in \{CDP_{3,3,3}, CEP_{T,T,T}, CP_7P_{T,T,T}, CB_{1,3}R\}$.

Proof. Let C, X, Y be a good triple. The graphs $P_{3,3,3}$ and $P_{T,T,T}$ are 2-connected, non-hamiltonian and claw-free and thus the set of CXY -free graphs contains neither $P_{3,3,3}$ nor $P_{T,T,T}$. Consequently, the pair X, Y must contain both an induced subgraph of $P_{3,3,3}$ and an induced subgraph of $P_{T,T,T}$. All connected induced subgraphs of the graph $P_{3,3,3}$ which are not induced subgraphs of P_6 , N or W are listed in Fig. 3 and, analogously, all connected induced subgraphs of the graph $P_{T,T,T}$ which are not induced subgraphs of P_6 , N or W are listed in Fig. 4.

From these lists we can see that $P_{3,3,3}$ and $P_{T,T,T}$ contain no common induced subgraph which is not an induced subgraph of P_6 , N or W . This implies that if CXY is a good triple, then (up to symmetry) X must be a graph from Fig. 3 and Y a graph from Fig. 4. Thus at the moment we have 160 candidates for good triples.

Consider a graph $P_{i,j,k}$ such that $i, j, k \geq 4$. This graph is 2-connected, non-hamiltonian and claw-free. Moreover it contains no induced subgraph isomorphic to C_4 , C_6 , H , to two triangles joined by an edge, or B_4 . Thus, at least one of the graphs X and Y does




Fig. 4.

not contain a copy of these graphs as an induced subgraph. Among the graphs in Fig. 4 only A_8 (the deer) and among the graphs in Fig. 3 only the graphs B_6 , B_7 , B_8 and B_9 satisfy this condition.

Let $X = A_8$. Then the triple CA_8B_1 is in fact the triple $CDP_{3,3,3}$ and, by Theorem B, this is a good triple. It is easy to see that this triple is also maximal. From this it immediately follows that all triples CA_8B_i where $i \neq 1$ are good but not maximal. Thus, it remains to consider all pairs $X = A_i, Y = B_j$ such that $i \neq 8$ and $j = 6, 7, 8, 9$.

Now the triple CA_1B_7 is in fact the triple $CEP_{T,T,T}$, the triple CA_1B_8 is the triple $CP_7P_{T,T,T}$ and both these triples are good by Theorem B. It is easy to see that both these triples are also maximal. Thus, all the triples CA_iB_7 and CA_iB_8 for $i \neq 1, 8$ are good but not maximal.

It remains to consider the triples CA_iB_6 and CA_iB_9 for $i \neq 8$. In [5] it was proved that the only 2-connected non-hamiltonian CZ_3 -free graphs are the graphs $P_{T,T,T}$ and $P_{T,T,3}$ and it is easy to see that these two graphs are also the only two graphs from the class \mathcal{P} which are B_6 -free. If we remove a vertex of degree four from $P_{T,T,T}$ and a vertex of degree three from $P_{T,T,3}$ then in both these cases we get the graph A_3 (which is isomorphic to the racket R). This implies that the triple $CB_{1,3}R$ is a good triple. Hence, all triples CA_iB_j where $j = 6, 9$ and A_i is an induced subgraph of $A_3 = R$ are good but not maximal. These are all graphs from Fig. 4 except A_1 , A_2 , A_4 and A_9 .

Now we have last eight triples CA_iB_j for $i = 1, 2, 4, 9$ and $j = 6, 9$. But the graph $P_{T,T,3}$ contains no such graph as an induced subgraph. Thus, these triples are not good triples and consequently the triple CA_3B_6 is a maximal good triple. The proof is complete. \square

Acknowledgements

The main result of this paper was achieved in the stimulating atmosphere of the 5th workshop “Cycles and Colourings” organized at Stará Lesná by Department of Geometry and Algebra, UJPŠ Košice, Slovakia, September 1998.

References

- [1] P. Bedrossian, Forbidden subgraphs and minimum degree conditions for hamiltonicity, Ph.D. Thesis, Memphis State University, USA, 1991.
- [2] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, Macmillan, London, Elsevier, New York, 1976.
- [3] J. Brousek, Minimal 2-connected non-hamiltonian claw-free graphs, Discrete Math. 191 (1998) 57–64.
- [4] R.J. Faudree, R.J. Gould, M. Jacobson, L. Lesniak, Triples of forbidden subgraphs implying hamiltonicity, preprint 1998.
- [5] R.J. Faudree, R.J. Gould, Z. Ryjáček, I. Schiermeyer, Forbidden subgraphs and pancyclicity, Congr. Numer. 109 (1995) 13–32.